

Navier-Stokes evolutions as selfdual variational problems

Nassif GHOUSSOUB
Department of Mathematics
University of British Columbia
Annex, Rm 1214
Vancouver, BC V6T 1Z2
CANADA

nassif@math.ubc.ca

Abstract

Many nonlinear PDEs and evolutions—including stationary and dynamic Navier-Stokes equations—can be formulated and resolved variationally by minimizing energy functionals of the form

$$I(u) = L(u, -\Lambda u) + \langle \Lambda u, u \rangle$$

and

$$I(u) = \int_0^T \left[L(t, u(t), -\dot{u}(t) - \Lambda u(t)) + \langle \Lambda u(t), u(t) \rangle \right] dt \\ + \ell(u(0) - u(T), \frac{u(T) + u(0)}{2})$$

where L is a time-dependent “selfdual Lagrangian” on state space, ℓ is another selfdual “boundary Lagrangian”, and Λ is a nonlinear operator (such as $\Lambda u = \operatorname{div}(u \otimes u)$ in the NSE case). However, just like the selfdual Yang–Mills equations, the equations are not obtained via Euler–Lagrange theory, but from the fact that a natural infimum is attained. In dimension 2, we recover the well known solutions for the corresponding initial-value problem as well as periodic and anti-periodic ones, while in dimension 3 we get Leray solutions for the initial-value problems, but also solutions satisfying $u(0) = \alpha u(T)$ for any given α in $(-1, 1)$. It is worth noting that our variational principles translate into Leray’s energy identity in dimension 2 (resp., inequality in dimension 3). Our approach is quite general and does apply to many other situations.