

Conservative Partially Hyperbolic Dynamics

Amie Wilkinson*

Abstract

We discuss recent progress in understanding the dynamical properties of partially hyperbolic diffeomorphisms that preserve volume. The main topics addressed are density of stable ergodicity and stable accessibility, center Lyapunov exponents, pathological foliations, rigidity, and the surprising interrelationships between these notions.

Mathematics Subject Classification (2000). Primary 37D30; Secondary 37C40.

Introduction

Here is a story, told at least in part through the exploits of one of its main characters. This character, like many a Hollywood (or Bollywood) star, has played a leading role in quite a few compelling tales; this one ultimately concerns the dynamics of partially hyperbolic diffeomorphisms.

We begin with a connected, compact, smooth surface S without boundary, of genus at least 2. The Gauss-Bonnet theorem tells us that the average curvature of any Riemannian metric on S must be negative, equal to $2\pi\chi(S)$, where $\chi(S)$ is the Euler characteristic of S . We restrict our attention to the metrics on S of everywhere negative curvature; among such metrics, there is a finite-dimensional moduli space of *hyperbolic* metrics, which have constant curvature. Up to a normalization of the curvature, each hyperbolic surface may be represented by a quotient \mathbb{H}/Γ , where \mathbb{H} is the complex upper half plane with the metric $y^{-2}(dx^2 + dy^2)$, and Γ is a discrete subgroup of $\mathrm{PSL}(2, \mathbb{R})$, isomorphic to the fundamental group of S . More generally, any negatively curved metric

*Thanks to Christian Bonatti, Keith Burns, Jordan Ellenberg, Andy Hammerlindl, François Ledrappier, Charles Pugh, Mike Shub and Lee Wilkinson for reading earlier versions of this text and making several helpful suggestions. This work was supported by the NSF.

Department of Mathematics, Northwestern University, 2033 Sheridan Road, Evanston, IL 60208-2730, USA. E-mail: wilkinso@math.northwestern.edu.

on S lies in the conformal class of some hyperbolic metric, and the space of all such metrics is path connected. Throughout this story, S will be equipped with a negatively curved metric.

This negatively curved muse first caught the fancy of Jacques Hadamard in the late 1890's [39]. Among other things, Hadamard studied the properties of geodesics on S and a flow $\varphi_t: T^1S \rightarrow T^1S$ on the unit tangent bundle to S called the *geodesic flow*. The image of a unit vector v under the time- t map of this flow is obtained by following the unique unit-speed geodesic $\gamma_v: \mathbb{R} \rightarrow S$ satisfying $\dot{\gamma}_v(0) = v$ for a distance t and taking the tangent vector at that point:

$$\varphi_t(v) := \dot{\gamma}_v(t).$$

This geodesic flow, together with its close relatives, plays the starring role in the story told here.




Figure 1. The geodesic flow.

A theorem of Liouville implies that φ_t preserves a natural probability measure m on T^1S , known as *Liouville measure*, which locally is just the product of normalized area on S with Lebesgue measure on the circle fibers. Poincaré recurrence then implies that almost every orbit of the geodesic flow comes back close to itself infinitely often.

In the special case where $S = \mathbb{H}/\Gamma$ is a hyperbolic surface, the unit tangent bundle T^1S is naturally identified with $PSL(2, \mathbb{R})/\Gamma$, and the action of the geodesic flow φ_t is realized by left multiplication by the diagonal matrix

$$g_t = \begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix}.$$

Liouville measure is normalized Haar measure.

In his study of φ_t , Hadamard introduced the notion of the *stable manifold* of a vector $v \in T^1S$:

$$\mathcal{W}^s(v) := \left\{ w \in T^1S \mid \lim_{t \rightarrow \infty} \text{dist}(\varphi_t(v), \varphi_t(w)) = 0 \right\}.$$

The proof that such sets are manifolds is a nontrivial consequence of negative curvature and a noted accomplishment of Hadamard's. Indeed, each stable

manifold $\mathcal{W}^s(v)$ is an injectively immersed, smooth copy of the real line, and taken together, the stable manifolds form a foliation \mathcal{W}^s of T^1M . Similarly, one defines an *unstable manifold* by:

$$\mathcal{W}^u(v) := \left\{ w \in T^1S \mid \lim_{t \rightarrow -\infty} \text{dist}(\varphi_t(v), \varphi_t(w)) = 0 \right\}$$

and denotes the corresponding unstable foliation \mathcal{W}^u . The foliations \mathcal{W}^s and \mathcal{W}^u are key supporting players in this story.

In the case where $S = \mathbb{H}/\Gamma$, the stable manifolds are orbits of the *positive horocyclic flow* on $\text{PSL}(2, \mathbb{R})/\Gamma$ defined by left-multiplication by

$$h_t^s = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix},$$

and the unstable manifolds are orbits of the *negative horocyclic flow*, defined by left-multiplication by

$$h_t^u = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}.$$

This fact can be deduced from the explicit relations:

$$g_{-t} h_r^s g_t = h_{re^{-t}}^s \quad \text{and} \quad g_{-t} h_r^u g_t = h_{re^t}^u. \quad (1)$$

The stable and unstable foliations stratify the future and past, respectively, of the geodesic flow. It might come as no surprise that their features dictate the asymptotic behavior of the geodesic flow. For example, Hadamard obtained from the existence of these foliations and Poincaré recurrence that periodic orbits for φ_t are dense in T^1S .

Some 40 years after Hadamard received the Prix Poncelet for his work on surfaces, Eberhard Hopf introduced a simple argument that proved the ergodicity (with respect to Liouville measure) of the geodesic flow on T^1S , for any closed negatively curved surface S [44]. In particular, Hopf proved that almost every infinite geodesic in S is dense (and uniformly distributed), not only in S , but in T^1S . It was another thirty years before Hopf's result was extended by Anosov to geodesic flows for negatively curved compact manifolds in arbitrary dimension.

Up to this point the discussion is quite well-known and classical, and from here the story can take many turns. For example, for arithmetic hyperbolic surfaces, the distribution of closed orbits of the flow and associated dynamical zeta functions quickly leads us into deep questions in analytic number theory. Another path leads to the study the spectral theory of negatively curved surfaces, inverse problems and quantum unique ergodicity. The path we shall take here leads to the definition of partial hyperbolicity.

Let us fix a unit of time $t_0 > 0$ and discretize the system φ_t in these units; that is, we study the dynamics of the time- t_0 map φ_{t_0} of the geodesic flow. From a digital age perspective this is a natural thing to do; for example, to plot

the orbits of a flow, a computer evaluates the flow at discrete, usually equal, time intervals.

If we carry this computer-based analogy one step further, we discover an interesting question. Namely, a computer does not “evaluate the flow” precisely, but rather uses an *approximation to the time- t_0 map* (such as an ODE solver or symplectic integrator) to compute its orbits. To what extent does iterating this approximation retain the actual dynamical features of the flow φ_t , such as ergodicity?

To formalize this question, we consider a *diffeomorphism* $f: T^1S \rightarrow T^1S$ such that the C^1 distance $d_{C^1}(f, \varphi_{t_0})$ is small. Note that f in general will no longer embed in a flow. While we assume that the distance from f to φ_{t_0} is small, this is no longer the case for the distance from f^n to φ_{nt_0} , when n is large.




Figure 2. $f^n(x)$ is not a good approximation to $\varphi_{nt_0}(x)$.

The earliest description of the dynamics of such a perturbation f comes from a type of structural stability theorem proved by Hirsch, Pugh, and Shub [43]. The results there imply in particular that if $d_{C^1}(f, \varphi_{t_0})$ is sufficiently small, then there exists an f -invariant *center foliation* $\mathcal{W}^c = \mathcal{W}^c(f)$ that is homeomorphic to the orbit foliation \mathcal{O} of φ_t . The leaves of \mathcal{W}^c are smooth. Moreover, the homeomorphism $h: T^1S \rightarrow T^1S$ sending \mathcal{W}^c to \mathcal{O} is close to the identity and \mathcal{W}^c is the unique such foliation.

The rest of this paper is about f and, in places, the foliation $\mathcal{W}^c(f)$.

What is known about f is now substantial, but far from complete. For example, the following basic problem is open.

Problem. Determine whether f has a dense orbit. More precisely, does there exist a neighborhood \mathcal{U} of φ_{t_0} in the space $\text{Diff}^r(T^1S)$ of C^r diffeomorphisms of T^1S (for some $r \geq 1$) such that every $f \in \mathcal{U}$ is topologically transitive?

Note that φ_{t_0} is ergodic with respect to volume m , and hence is itself topologically transitive. In what follows, we will explain results from the last 15 years implying that any perturbation of φ_{t_0} that preserves volume is ergodic, and hence has a dense orbit. For perturbations that do not preserve volume, a seminal result of Bonatti and Díaz shows that φ_{t_0} can be approximated arbitrarily well by C^1 -open sets of transitive diffeomorphisms [9]. But the fundamental question of whether φ_{t_0} lives in such an open set remains unanswered.

In most of the discussion here, we will work in the conservative setting, in which the diffeomorphism f preserves a volume probability measure. To fix notation, M will always denote a connected, compact Riemannian manifold without boundary, and m will denote a probability volume on M . For $r \geq 1$, we denote by $\text{Diff}_m^r(M)$ the space of C^r diffeomorphisms of M preserving m , equipped with the C^r topology.

1. Partial Hyperbolicity

The map φ_{t_0} and its perturbation f are concrete examples of partially hyperbolic diffeomorphisms. A diffeomorphism $f: M \rightarrow M$ of a compact Riemannian manifold M is *partially hyperbolic* if there exists an integer $k \geq 1$ and a non-trivial, Df -invariant, continuous splitting of the tangent bundle

$$TM = E^s \oplus E^c \oplus E^u$$

such that, for any $p \in M$ and unit vectors $v^s \in E^s(p)$, $v^c \in E^c(p)$, and $v^u \in E^u(p)$:

$$\begin{aligned} \|D_p f^k v^s\| &< 1 < \|D_p f^k v^u\|, & \text{and} \\ \|D_p f^k v^s\| &< \|D_p f^k v^c\| < \|D_p f^k v^u\|. \end{aligned}$$

Up to a change in the Riemannian metric, one can always take $k = 1$ in this definition [37]. In the case where E^c is the trivial bundle, the map f is said to be *Anosov*. The central example φ_{t_0} is partially hyperbolic: in that case, the bundle $E^c = \mathbb{R}\dot{\varphi}$ is tangent to the orbits of the flow, and E^s and E^u are tangent to the leaves of \mathcal{W}^s and \mathcal{W}^u , respectively.

Partial hyperbolicity is a C^1 -open condition: any diffeomorphism sufficiently C^1 -close to a partially hyperbolic diffeomorphism is itself partially hyperbolic. Hence the perturbations of φ_{t_0} we consider are also partially hyperbolic. For an extensive discussion of examples of partially hyperbolic dynamical systems, see the survey articles [20, 41, 62] and the book [55]. Among these examples are: the frame flow for a compact manifold of negative sectional curvature and most affine transformations of compact homogeneous spaces.

As is the case with the example φ_{t_0} , the stable and unstable bundles E^s and E^u of an arbitrary partially hyperbolic diffeomorphism are always tangent to foliations, which we will again denote by \mathcal{W}^s and \mathcal{W}^u respectively; this is a consequence of partial hyperbolicity and a generalization of Hadamard's argument. By contrast, the center bundle E^c need not be tangent to a foliation,

and can even be nowhere integrable. In many cases of interest, however, there is also a center foliation \mathcal{W}^c tangent to E^c : the content of the Hirsch-Pugh-Shub work in [43] is the properties of systems that admit such foliations, known as “normally hyperbolic foliations.”

There is a natural and slightly less general notion than integrability of E^c that appears frequently in the literature. We say that a partially hyperbolic diffeomorphism $f: M \rightarrow M$ is *dynamically coherent* if the subbundles $E^c \oplus E^s$ and $E^c \oplus E^u$ are tangent to foliations \mathcal{W}^{cs} and \mathcal{W}^{cu} , respectively, of M . If f is dynamically coherent, then the center bundle E^c is also integrable: one obtains the center foliation \mathcal{W}^c by intersecting the leaves of \mathcal{W}^{cs} and \mathcal{W}^{cu} . The examples φ_{t_0} are dynamically coherent, as are their perturbations (by [43]: see [24] for a discussion).

2. Stable Ergodicity and the Pugh-Shub Conjectures

Brin and Pesin [16] and independently Pugh and Shub [57] first examined the ergodic properties of partially hyperbolic systems in the early 1970’s. The methods they developed give an ergodicity criterion for partially hyperbolic $f \in \text{Diff}_m^2(M)$ satisfying the following additional hypotheses:

- (a) the bundle E^c is tangent to a C^1 foliation \mathcal{W}^c , and
- (b) f acts isometrically (or nearly isometrically) on the leaves of \mathcal{W}^c .

In [16] it is shown that such an f is ergodic with respect to m if it satisfies a condition called accessibility.

Definition 2.1. A partially hyperbolic diffeomorphism $f: M \rightarrow M$ is *accessible* if any point in M can be reached from any other along an *su-path*, which is a concatenation of finitely many subpaths, each of which lies entirely in a single leaf of \mathcal{W}^s or a single leaf of \mathcal{W}^u .

This ergodicity criterion applies to the discretized geodesic flow φ_{t_0} : its center bundle is tangent to the orbit foliation for φ_t , which is smooth, giving

- (a). The action of φ_{t_0} preserves the nonsingular vector field $\dot{\varphi}$, which implies
- (b). It is straightforward to see that if S is a hyperbolic surface, then φ_{t_0} is accessible: the stable and unstable foliations are orbits of the smooth horocyclic flows h_t^s and h_t^u , respectively, and matrix multiplication on the level of the Lie algebra \mathfrak{sl}_2 shows that locally these flows generate all directions in $\text{PSL}(2, \mathbb{R})$:

$$\frac{1}{2} \left[\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right] = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}; \quad (2)$$

the matrices appearing on the left are infinitesimal generators of the horocyclic flows, and the matrix on the right generates the geodesic flow. Since φ_{t_0} is accessible, it is ergodic.

Now what of a small perturbation of φ_{t_0} ? As mentioned above, any $f \in \text{Diff}_m^2(T^1S)$ sufficiently C^1 close to φ_{t_0} also has a center foliation \mathcal{W}^c , and the action of f on the leaves is nearly isometric. With some work, one can also show that f is accessible (this was carried out in [16]). There is one serious reason why the ergodicity criterion of [16] cannot be applied to f : the foliation \mathcal{W}^c is not C^1 . The leaves of \mathcal{W}^c are C^1 , and the tangent spaces to the leaves vary continuously, but they do not vary smoothly. We will explore in later sections the extent to which \mathcal{W}^c fails to be smooth, but for now suffice it to say that \mathcal{W}^c is pathologically bad, not only from a smooth perspective but also from a measure-theoretic one.

The extent to which \mathcal{W}^c is bad was not known at the time, but there was little hope of applying the existing techniques to perturbations of φ_{t_0} . The first major breakthrough in understanding the ergodicity of perturbations of φ_{t_0} came in the 1990's:

Theorem A (Grayson-Pugh-Shub [38]). *Let S be a hyperbolic surface, and let φ_t be the geodesic flow on T^1S . Then φ_{t_0} is stably ergodic: there is a neighborhood \mathcal{U} of φ_{t_0} in $\text{Diff}_m^2(T^1S)$ such that every $f \in \mathcal{U}$ is ergodic with respect to m .*

The new technique introduced in [38] was a dynamical approach to understanding Lebesgue density points which they called *juliennes*. The results in [38] were soon generalized to the case where S has variable negative curvature [74] and to more general classes of partially hyperbolic diffeomorphisms [59, 60]. Not long after [38] appeared, Pugh and Shub had formulated an influential circle of conjectures concerning the ergodicity of partially hyperbolic systems.

Conjecture 1 (Pugh-Shub [58]). *On any compact manifold, ergodicity holds for an open and dense set of C^2 volume preserving partially hyperbolic diffeomorphisms.*

This conjecture can be split into two parts using the concept of accessibility.

Conjecture 2 (Pugh-Shub [58]). *Accessibility holds for an open and dense subset of C^2 partially hyperbolic diffeomorphisms, volume preserving or not.*

Conjecture 3 (Pugh-Shub [58]). *A partially hyperbolic C^2 volume preserving diffeomorphism with the essential accessibility property is ergodic.*

Essential accessibility is a measure-theoretic version of accessibility that is implied by accessibility: f is essentially accessible if for any two positive volume sets A and B , there exists an su -path in M connecting some point in A to some point in B – see [20] for a discussion.

In the next two sections, I will report on progress to date on these conjectures.

Further remarks.

1. Volume-preserving Anosov diffeomorphisms (where $\dim E^c = 0$) are always ergodic. This was proved by Anosov in his thesis [1]. Note that Anosov diffeomorphisms are also accessible, since in that case the foliations \mathcal{W}^s and \mathcal{W}^u are transverse. Hence all three conjectures hold true for Anosov diffeomorphisms.
2. It is natural to ask whether partial hyperbolicity is a necessary condition for stable ergodicity. This is true when M is 3-dimensional [27] and also in the space of symplectomorphisms [45, 68], but not in general [71]. What is true is that the related condition of having a dominated splitting is necessary for stable ergodicity (see [27]).
3. One can also ask whether for partially hyperbolic systems, stable ergodicity implies accessibility. If one works in a sufficiently high smoothness class, then this is not the case, as was shown in the groundbreaking paper of F. Rodríguez Hertz [61], who will also speak at this congress. Hertz used methods from KAM theory to find an alternate route to stable ergodicity for certain essentially accessible systems.
4. On the other hand, it is reasonable to expect that some form of accessibility is a necessary hypothesis for a general stable ergodicity criterion for partially hyperbolic maps (see the discussion at the beginning of [24]). Unlike Anosov diffeomorphisms, which are always ergodic, partially hyperbolic diffeomorphisms need not be ergodic. For example, the product of an Anosov diffeomorphism with the identity map on any manifold is partially hyperbolic, but certainly not ergodic. See also Theorem 11.16 in [10].

3. Accessibility

In general, the stable and unstable foliations of a partially hyperbolic diffeomorphism are not smooth (though they are not pathological, either – see below). Hence it is not possible in general to use infinitesimal techniques to establish accessibility the way we did in equation (2) for the discretized hyperbolic geodesic flow. The C^1 topology allows for enough flexibility in perturbations that Conjecture 2 has been completely verified in this context:

Theorem B (Dolgopyat-Wilkinson [31]). *For any $r \geq 1$, accessibility holds for a C^1 open and dense subset of the partially hyperbolic diffeomorphisms in $\text{Diff}^r(M)$, volume-preserving or not.*

Theorem B also applies inside the space of partially hyperbolic symplectomorphisms.

More recently, the complete version of Conjecture 3 has been verified for systems with 1-dimensional center bundle.

Theorem C (Rodríguez Hertz-Rodríguez Hertz-Ures [63]). *For any $r \geq 1$, accessibility is C^1 open and C^r dense among the partially hyperbolic diffeomorphisms in $\text{Diff}_m^r(M)$ with one-dimensional center bundle.*

This theorem was proved earlier in a much more restricted context by Nițică-Török [54]. The C^1 openness of accessibility was shown in [28]. A version of Theorem C for non-volume preserving diffeomorphisms was later proved in [19].

The reason that it is possible to improve Theorem B from C^1 density to C^r density in this context is that the global structure of accessibility classes is well-understood. By *accessibility class* we mean an equivalence class with respect to the relation generated by *su*-paths. When the dimension of E^c is 1, accessibility classes are (C^1 immersed) submanifolds. Whether this is always true when $\dim(E^c) > 1$ is unknown and is an important obstacle to attacking the general case of Conjecture 2.

Further remarks.

1. More precise criteria for accessibility have been established for special classes of partially hyperbolic systems such as discretized Anosov flows, skew products, and low-dimensional systems [23, 21, 64].
2. Refined formulations of accessibility have been used to study higher-order statistical properties of certain partially hyperbolic systems, in particular the discretized geodesic flow [29, 52]. The precise relationship between accessibility and rate of mixing (in the absence of other hypotheses) remains a challenging problem to understand.
3. Accessibility also plays a key role in a recently developed Livsič theory for partially hyperbolic diffeomorphisms, whose conclusions closely mirror those in the Anosov setting [47, 73].

4. Ergodicity

Conjecture 1 has been verified under one additional, reasonably mild hypothesis:

Theorem D (Burns-Wilkinson [22]). *Let f be C^2 , volume-preserving, partially hyperbolic and center bunched. If f is essentially accessible, then f is ergodic, and in fact has the Kolmogorov property.*

The additional hypothesis is “center bunched.” A partially hyperbolic diffeomorphism f is *center bunched* if there exists an integer $k \geq 1$ such that for any $p \in M$ and any unit vectors $v^s \in E^s(p)$, $v^c, w^c \in E^c(p)$, and $v^u \in E^u(p)$:

$$\|D_p f^k v^s\| \cdot \|D_p f^k w^c\| < \|D_p f^k v^c\| < \|D_p f^k v^u\| \cdot \|D_p f^k w^c\|. \quad (3)$$

As with partial hyperbolicity, the definition of center bunching depends only on the smooth structure on M and not the Riemannian structure; if (3) holds for a

given metric and $k \geq 1$, one can always find another metric for which (3) holds with $k = 1$ [37]. In words, center bunching requires that the non-conformality of $Df|_{E^c}$ be dominated by the hyperbolicity of $Df|_{E^u \oplus E^s}$. Center bunching holds automatically if the restriction of Df to E^c is conformal in some metric (for this metric, one can choose $k = 1$). In particular, if E^c is one-dimensional, then f is center bunched. In the context where $\dim(E^c) = 1$, Theorem D was also shown in [63].

Combining Theorems C and D we obtain:

Corollary 1. *The Pugh-Shub conjectures hold true among the partially hyperbolic diffeomorphisms with 1-dimensional center bundle.*

Further remarks.

1. The proof of Theorem D builds on the original argument of Hopf for ergodicity of geodesic flows and incorporates a refined theory of the julienne originally introduced in [38].
2. It appears that the center bunching hypothesis in Theorem D cannot be removed without a significantly new approach. On the other hand, it is possible that Conjecture 1 will yield to other methods.
3. Formulations of Conjecture 1 in the C^1 topology have been proved for low-dimensional center bundle [12, 61] and for symplectomorphisms [3]. These formulations state that ergodicity holds for a *residual* subset in the C^1 topology.

5. Exponents

By definition, a partially hyperbolic diffeomorphism produces uniform contraction and expansion in the directions tangent to E^s and E^u , respectively. In none of the results stated so far do we make any precise assumption on the growth of vectors in E^c beyond the coarse bounds that come from partial hyperbolicity and center bunching. In particular, an ergodic diffeomorphism in Theorem D can have periodic points of different indices, corresponding to places in M where E^c is uniformly expanded, contracted, or neither. The power of the julienne-based theory is that the hyperbolicity in $E^u \oplus E^s$, when combined with center bunching and accessibility, is enough to cause substantial mixing in the system, regardless of the precise features of the dynamics on E^c .

On the other hand, the asymptotic expansion/contraction rates in E^c can give additional information about the dynamics of the diffeomorphism, and is a potentially important tool for understanding partially hyperbolic diffeomorphisms that are not center bunched.

A real number λ is a *center Lyapunov exponent* of the partially hyperbolic diffeomorphism $f : M \rightarrow M$ if there exists a nonzero vector $v \in E^c$ such that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|Df^n(v)\| = \lambda. \quad (4)$$

If f preserves m , then Oseledec's theorem implies that the limit in (4) exists for each $v \in E^c(x)$, for m -almost every x . When the dimension of E^c is 1, the limit in (4) depends only on x , and if in addition f is ergodic with respect to m , then the limit takes a single value m -almost everywhere.

Theorem E (Shub-Wilkinson [70]). *There is an open set $\mathcal{U} \subset \text{Diff}_m^\infty(\mathbb{T}^3)$ of partially hyperbolic, dynamically coherent diffeomorphisms of the 3-torus $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$ for which:*

- *the elements of \mathcal{U} approximate arbitrarily well (in the C^∞ topology) the linear automorphism of \mathbb{T}^3 induced by the matrix:*

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- *the elements of \mathcal{U} are ergodic and have positive center exponents, m -almost everywhere.*

Note that the original automorphism A has vanishing center exponents, *everywhere* on \mathbb{T}^3 , since A is the identity map on the third factor. Yet Theorem E says that a small perturbation mixing the unstable and center directions of A creates expansion in the center direction, almost everywhere on \mathbb{T}^3 .

The systems in \mathcal{U} enjoy the feature of being *non-uniformly hyperbolic*: the Lyapunov exponents in every direction (not just center ones) are nonzero, m -almost everywhere. The well-developed machinery of Pesin theory guarantees a certain level of chaotic behavior from nonuniform hyperbolicity alone. For example, a nonuniformly hyperbolic diffeomorphism has at most countably many ergodic components, and a mixing partially hyperbolic diffeomorphism is Bernoulli (i.e. abstractly isomorphic to a Bernoulli process). A corollary of Theorem E is that the elements of \mathcal{U} are Bernoulli systems.

The constructions in [70] raise the question of whether it might be possible to “remove zero exponents” from any partially hyperbolic diffeomorphism via a small perturbation. If so, then one might be able to bypass the julienne based theory entirely and use techniques from Pesin theory instead as an approach to Conjecture 1. More generally, and wildly optimistically, one might ask whether *any* $f \in \text{Diff}_m^2(M)$ with at least one nonzero Lyapunov exponent on a positive measure set might be perturbed to produce nonuniform hyperbolicity on a positive measure set (such possibilities are discussed in [70]).

There is a partial answer to these questions in the C^1 topology, due to Baraveira and Bonatti [7]. The results there imply in particular that if $f \in$

$\text{Diff}_m^r(M)$ is partially hyperbolic, then there exists $g \in \text{Diff}_m^r(M)$, C^1 -close to f so that the *sum* of the center Lyapunov exponents is nonzero.

Further remarks.

1. Dolgopyat proved that the same type of construction as in [70] can be applied to the discretized geodesic flow φ_{t_0} for a negatively curved surface S to produce perturbations with nonzero center exponents [30]. See also [66] for further generalizations of [70].
2. An alternate approach to proving Conjecture 1 has been proposed, taking into account the center Lyapunov exponents [18]. For systems with $\dim(E^c) = 2$, this program has very recently been carried out in the C^1 topology in [65], using a novel application of the technique of blenders, a concept introduced in [9].

6. Pathology

There is a curious by-product of nonvanishing Lyapunov exponents for the open set \mathcal{U} of examples in Theorem E. By [43], there is a center foliation \mathcal{W}^c for each $f \in \mathcal{U}$, homeomorphic to the trivial \mathbb{R}/\mathbb{Z} fibration of $\mathbb{T}^3 = \mathbb{T}^2 \times \mathbb{R}/\mathbb{Z}$; in particular, the center leaves are all compact. The almost everywhere exponential growth associated with nonzero center exponents is incompatible with the compactness of the center foliation, and so the full volume set with positive center exponent must meet almost every leaf in a zero set (in fact a finite set [67]).

The same type of phenomenon occurs in perturbations of the discretized geodesic flow φ_{t_0} . While in that case the leaves of \mathcal{W}^c are mostly noncompact, they are in a sense “dynamically compact.” An adaptation of the arguments in [67] shows that any perturbation of φ_{t_0} with nonvanishing center exponents, such as those constructed by Dolgopyat in [30], have atomic disintegration of volume along center leaves.

Definition 6.1. A foliation \mathcal{F} of M with smooth leaves has *atomic disintegration of volume* along its leaves if there exists $A \subset M$ such that

- $m(M \setminus A) = 0$, and
- A meets each leaf of \mathcal{F} in a discrete set of points (in the leaf topology).

At the opposite end of the spectrum from atomic disintegration of volume is a property called absolute continuity. A foliation \mathcal{F} is *absolutely continuous* if holonomy maps between smooth transversals send zero volume sets to zero volume sets. If \mathcal{F} has smooth leaves and is absolutely continuous, then for every set $A \subset M$ satisfying $m(M \setminus A) = 0$, the intersection of A with the leaf \mathcal{F} through m -almost every point in M has full leafwise Riemannian volume. In this sense Fubini’s theorem holds for absolutely continuous foliations. If \mathcal{F}




Figure 3. A pathological foliation

is a C^1 foliation, then it is absolutely continuous, but absolute continuity is a strictly weaker property.

Absolute continuity has long played a central role in smooth ergodic theory. Anosov and Sinai [1, 2] proved in the 60's that the stable and unstable foliations of globally hyperbolic (or Anosov) systems are absolutely continuous, even though they fail to be C^1 in general. Absolute continuity was a key ingredient in Anosov's celebrated proof [1] that the geodesic flow for any compact, negatively curved manifold is ergodic. When the center foliation for f fails to be absolutely continuous, this means that one cannot "quotient out by the center direction" to study ergodic properties f .

The existence of such pathological center foliations was first demonstrated by A. Katok (whose construction was written up by Milnor in [53]). Theorem E shows that this type of pathology can occur in open sets of diffeomorphisms and so is inescapable in general. In the next section, we discuss the extent to which this pathology is the norm.

Further remarks.

1. An unpublished letter of Mañé to Shub examines the consequences of nonvanishing Lyapunov center exponents on the disintegration of volume along center foliations. Some of the ideas there are pursued in greater depth in [42].
2. The examples of Katok in [53] in fact have center exponents almost everywhere equal to 0, showing that nonvanishing center exponents is not a necessary condition for atomic disintegration of volume.

3. Systems for which the center leaves are not compact (or even dynamically compact) also exhibit non-absolutely continuous center foliations, but the disintegration appears to be potentially much more complicated than just atomic disintegration [69, 36].

7. Rigidity

Examining in greater depth the potential pathologies of center foliations, we discover a rigidity phenomenon. To be concrete, let us consider the case of a perturbation $f \in \text{Diff}_m^\infty(M)$ of the discretized geodesic flow on a negatively-curved surface. If the perturbation f happens to be the time-one map of a smooth flow, then \mathcal{W}^c is the orbit foliation for that flow. In this case the center foliation for f is absolutely continuous – in fact, C^∞ . In general, however, a perturbation f of φ_{t_0} has no reason to embed in a smooth flow, and one can ask how the volume m disintegrates along the leaves of \mathcal{W}^c .

There is a complete answer to this question:

Theorem F (Avila-Viana-Wilkinson [6]). *Let S be a closed negatively curved surface, and let $\varphi_t : T^1S \rightarrow T^1S$ be the geodesic flow.*

For each $t_0 > 0$, there is a neighborhood \mathcal{U} of φ_{t_0} in $\text{Diff}_m^\infty(T^1S)$ such that for each $f \in \mathcal{U}$:

1. *either m has atomic disintegration along the center foliation \mathcal{W}^c , or*
2. *f is the time-one map of a C^∞ , m -preserving flow.*

What Theorem F says is that, in this context, nothing lies between C^∞ and absolute singularity of \mathcal{W}^c – *pathology is all that can happen*. The geometric measure-theoretic properties of \mathcal{W}^c determine completely a key dynamical property of f – whether it embeds in a flow.

The heart of the proof of Theorem F is to understand what happens when the center Lyapunov exponents *vanish*. For this, we use tools that originate in the study of random matrix products. The general theme of this work, summarized by Ledrappier in [49] is that “entropy is smaller than exponents, and entropy zero implies deterministic.” Original results concerning the Lyapunov exponents of random matrix products, due to Furstenberg, Kesten [34, 33], Ledrappier [49], and others, have been extended in the past decade to deterministic products of linear cocycles over hyperbolic systems by Bonatti, Gomez-Mont, Viana [11, 13, 72]. The Bernoulli and Markov measures associated with random products in those earlier works are replaced in the newer results by invariant measures for the hyperbolic system carrying a suitable product structure.

Recent work of Avila, Viana [5] extends this hyperbolic theory from linear to *diffeomorphism* cocycles, and these results are used in a central way. For cocycles over volume preserving partially hyperbolic systems, Avila, Santamaria,

and Viana [4] have also recently produced related results, for both linear and diffeomorphism cocycles, which also play an important role in the proof. The proof in [4] employs julienne based techniques, generalizing the arguments in [24].

Further remarks.

1. The only properties of φ_{t_0} that are used in the proof of Theorem F are accessibility, dynamical coherence, one-dimensionality of E^c , the fact that φ_{t_0} fixes the leaves of \mathcal{W}^c , and 3-dimensionality of M . There are also more general formulations of Theorem F in [6] that relax these hypotheses in various directions. For example, a similar result holds for systems in dimension 3 for whom all center manifolds are compact.
2. Deep connections between Lyapunov exponents and geometric properties of invariant measures have long been understood [48, 50, 51, 46, 8]. Theorem F establishes new connections in the partially hyperbolic context.
3. Theorem F gives conditions under which one can recover the action of a Lie group (in this case \mathbb{R}) from that of a discrete subgroup (in this case \mathbb{Z}). These themes have arisen in the related context of measure-rigidity for algebraic partially hyperbolic actions by Einsiedler, Katok, Lindenstrauss [32]. It would be interesting to understand more deeply the connections between these works.

8. Summary, Questions

We leave this tale open-ended, with a few questions that have arisen naturally in its course.

New criteria for ergodicity. Conjecture 1 remains open. As discussed in Section 4, the julienne based techniques using the Hopf argument might have reached their limits in this problem (at least this is the case in the absence of a significantly new idea). One alternate approach which seems promising employs Lyapunov exponents and blenders [65]. Perhaps a new approach will find a satisfying conclusion to this part of the story.

Classification problem. A basic question is to understand which manifolds support partially hyperbolic diffeomorphisms. As the problem remains open in the classical Anosov case (in which E^c is zero-dimensional), it is surely extremely difficult in general. There has been significant progress in dimension 3, however; for example, using techniques in the theory of codimension-1 foliations, Burago and Ivanov proved that there are no partially hyperbolic diffeomorphisms of the 3-sphere [17].

Modifying this question slightly, one can ask whether the partially hyperbolic diffeomorphisms in low dimension must belong to certain “classes” (up to

homotopy, for example) – such as time- t maps of flows, skew products, algebraic systems, and so on. Pujals has proposed such a program in dimension 3, which has spurred several papers on the subject [15, 14, 64, 40].

It is possible that if one adds the hypotheses of dynamical coherence and absolute continuity of the center foliation, then there is such a classification. Evidence in this direction can be found in [6].

Nonuniform and singular partial hyperbolicity. Unless all of its Lyapunov exponents vanish almost everywhere, *any* volume-preserving diffeomorphism is in some sense “nonuniformly partially hyperbolic.” Clearly such a general class of systems will not yield to a single approach. Nonetheless, the techniques developed recently are quite powerful and should shed some light on certain systems that are close to being partially hyperbolic. Some extensions beyond the uniform setting have been explored in [3], in which the center bunching hypotheses in [24] has been replaced by a pointwise, nonuniform center bunching condition. This gives new classes of stably ergodic diffeomorphisms that are not center bunched.

It is conceivable that the methods in [3] may be further extended to apply in certain “singular partially hyperbolic” contexts where partial hyperbolicity holds on an open, noncompact subset of the manifold M but decays in strength near the boundary. Such conditions hold, for example, for geodesic flows on certain nonpositively curved manifolds. Under suitable accessibility hypotheses, these systems should be ergodic with respect to volume.

Rigidity of partially hyperbolic actions. The rigidity phenomenon described in Section 7 has only begun to be understood. To phrase those results in a more general context, we consider a smooth, nonsingular action of an abelian Lie group G on a manifold M . Let H be a discrete group acting on M , commuting with the action of G , and whose elements are partially hyperbolic diffeomorphisms in $\text{Diff}_m^\infty(M)$. Can such an action be perturbed, preserving the absolute continuity of the center foliation? How about the elements of the action? When absolute continuity fails, what happens?

The role of accessibility and accessibility classes has been exploited in a serious way in the important work of Damjanović and A. Katok on rigidity of abelian actions on quotients of $\text{SL}(n, \mathbb{R})$ [26]. It seems reasonable that these explorations can be pushed further, using some of the techniques mentioned here, to prove rigidity results for other partially hyperbolic actions. A simple case currently beyond the reach of existing methods is to understand perturbations of the action of a \mathbb{Z}^2 lattice in the diagonal subgroup on $\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R})/\Gamma$, where Γ is an irreducible lattice.

Our final question takes us further afield, but back once again to the geodesic flow. Fix a closed hyperbolic surface S , and consider the standard action on T^1S by the upper triangular subgroup $T < \text{PSL}(2, \mathbb{R})$, which contains both the geodesic and positive horocyclic flows. Ghys proved that this action is highly rigid and admits no m -preserving C^∞ deformations [35]. Does the same hold true for some countable subgroup of T ? For example, consider the solvable

Baumslag Solitar subgroup $BS(1, 2)$ generated by the elements

$$a = \begin{pmatrix} \sqrt{2} & 0 \\ 0 & \frac{1}{\sqrt{2}} \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

which has the presentation $BS(1, 2) = \langle a, b \mid aba^{-1} = b^2 \rangle$. Can the standard action be perturbed inside of $\text{Diff}_m^\infty(T^1S)$? More generally, can one classify all faithful representations

$$\rho: BS(1, 2) \rightarrow \text{Diff}_m^\infty(M),$$

where M is a 3-manifold? For results of a similar nature in lower dimensions, see [25, 56].

References

- [1] D. V. Anosov, *Geodesic flows on closed Riemannian manifolds of negative curvature*, Proc. Steklov Math. Inst. **90** (1967), 1–235.
- [2] D. V. Anosov and Ya. G. Sinai, *Certain smooth ergodic systems*, Russian Math. Surveys **22** (1967), 103–167.
- [3] A. Avila, J. Bochi, and A. Wilkinson, *Nonuniform center bunching and the genericity of ergodicity among C^1 partially hyperbolic symplectomorphisms*.
- [4] A. Avila, J. Santamaria, and M. Viana, *Cocycles over partially hyperbolic maps*, Preprint www.preprint.impa.br 2008.
- [5] A. Avila and M. Viana, *Extremal Lyapunov exponents of smooth cocycles*, Invent. Math.
- [6] A. Avila, M. Viana, and A. Wilkinson, *Absolute continuity, Lyapunov exponents, and rigidity*, in preparation.
- [7] A. Baraviera and C. Bonatti, *Removing zero central Lyapunov exponents*, Ergod. Th. & Dynam. Sys. **23** (2003), 1655–1670.
- [8] L. Barreira, Ya. Pesin, and J. Schmeling, *Dimension and product structure of hyperbolic measures*, Ann. of Math. **149** (1999), 755–783.
- [9] C. Bonatti and L. J. Díaz, *Nonhyperbolic transitive diffeomorphisms*, Ann. of Math. **143** (1996), 357–396.
- [10] C. Bonatti, L. J. Díaz, and M. Viana, *Dynamics beyond uniform hyperbolicity*, Encyclopaedia of Mathematical Sciences, vol. 102, Springer-Verlag, 2005.
- [11] C. Bonatti, X. Gómez-Mont, and M. Viana, *Généricité d'exposants de Lyapunov non-nuls pour des produits déterministes de matrices*, Ann. Inst. H. Poincaré Anal. Non Linéaire **20** (2003), 579–624.
- [12] C. Bonatti, C. Matheus, M. Viana, and A. Wilkinson, *Abundance of stable ergodicity*, Comment. Math. Helv. **79** (2004), 753–757.

-
- [13] C. Bonatti and M. Viana, *Lyapunov exponents with multiplicity 1 for deterministic products of matrices*, Ergod. Th. & Dynam. Sys **24** (2004), 1295–1330.
- [14] C. Bonatti and A. Wilkinson, *Transitive partially hyperbolic diffeomorphisms on 3-manifolds*, Topology **44** (2005), 475–508.
- [15] M. Brin, D. Burago, and S. Ivanov, *On partially hyperbolic diffeomorphisms on 3-manifolds with commutative fundamental group*, Advances in Dynamical Systems, Cambridge Univ. Press.
- [16] M. Brin and Ya. Pesin, *Partially hyperbolic dynamical systems*, Izv. Acad. Nauk. SSSR **1** (1974), 177–212.
- [17] D. Burago and S. Ivanov, *Partially hyperbolic diffeomorphisms of 3-manifolds with abelian fundamental groups*, J. Mod. Dyn. **2** (2008), 541–580.
- [18] K. Burns, D. Dolgopyat, and Ya. Pesin, *Partial hyperbolicity, Lyapunov exponents and stable ergodicity*, J. Statist. Phys. **108** (2002), 927–942, Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.
- [19] K. Burns, F. Rodríguez Hertz, M. A. Rodríguez Hertz, A. Talitskaya, and R. Ures, *Density of accessibility for partially hyperbolic diffeomorphisms with one-dimensional center*, Discrete Contin. Dyn. Syst. **22** (2008), 75–88.
- [20] K. Burns, C. Pugh, M. Shub, and A. Wilkinson, *Recent results about stable ergodicity*, Smooth ergodic theory and its applications (Seattle WA, 1999), Procs. Symp. Pure Math., vol. 69, Amer. Math. Soc., 2001, pp. 327–366.
- [21] K. Burns, C. Pugh, and A. Wilkinson, *Stable ergodicity and Anosov flows*, Topology **39** (2000), 149–159.
- [22] K. Burns and A. Wilkinson, *On the ergodicity of partially hyperbolic systems*, Ann. of Math.
- [23] ———, *Stable ergodicity of skew products*, Ann. Sci. École Norm. Sup. **32** (1999), 859–889.
- [24] ———, *Dynamical coherence and center bunching*, Discrete Contin. Dyn. Syst. **22** (2008), 89–100.
- [25] L. Burslem and A. Wilkinson, *Global rigidity of solvable group actions on S^1* , Geometry and Topology **8** (2004), 877–924.
- [26] D. Damjanović and A. Katok, *Local rigidity of partially hyperbolic actions. II. The geometric method and restrictions of weyl chamber flows on $SL(n, \mathbb{R})/L$* .
- [27] L. J. Díaz, E. Pujals, and R. Ures, *Partial hyperbolicity and robust transitivity*, Acta Math. **183** (1999), 1–43.
- [28] P. Didier, *Stability of accessibility*, Ergod. Th. & Dynam. Sys. **23** (2003), 1717–1731.
- [29] D. Dolgopyat, *On decay of correlations in Anosov flows*, Ann. of Math. **147** (1998), 357–390.
- [30] ———, *On differentiability of SRB states for partially hyperbolic systems*, Invent. Math. **155** (2004), 389–449.
- [31] D. Dolgopyat and A. Wilkinson, *Stable accessibility is C^1 dense*, Astérisque **287** (2003), 33–60.

-
- [32] M. Einsiedler, A. Katok, and E. Lindenstrauss, *Invariant measures and the set of exceptions to Littlewood's conjecture*, Ann. of Math. **164** (2006), 513–560.
- [33] H. Furstenberg, *Non-commuting random products*, Trans. Amer. Math. Soc. **108** (1963), 377–428.
- [34] H. Furstenberg and H. Kesten, *Products of random matrices*, Ann. Math. Statist. **31** (1960), 457–469.
- [35] É. Ghys, *Actions localement libres du groupe affine*, Invent. Math. **82** (1985), 479–526.
- [36] A. Gogolev, *How typical are pathological foliations in partially hyperbolic dynamics: an example*, arXiv:0907.3533.
- [37] N. Gourmelon, *Adapted metrics for dominated splittings*, Ergod. Th. & Dynam. Sys. **27** (2007), 1839–1849.
- [38] M. Grayson, C. Pugh, and M. Shub, *Stably ergodic diffeomorphisms*, Ann. of Math. **140** (1994), 295–329.
- [39] J. Hadamard, *Les surfaces à courbures opposées et leurs lignes géodésiques*, Journal de Math. Pures et Appliquées **IV** (1898), 27–73.
- [40] A. Hammerlindl, *Leaf conjugacies on the torus*, Ph.D. thesis, U. Toronto, 2009.
- [41] B. Hasselblatt and Ya. Pesin, *Partially hyperbolic dynamical systems*, Handbook of dynamical systems. Vol. 1B, Elsevier B. V., 2006, pp. 1–55.
- [42] M. Hirayama and Ya. Pesin, *Non-absolutely continuous foliations*, Israel J. Math. **160** (2007), 173–187.
- [43] M. Hirsch, C. Pugh, and M. Shub, *Invariant manifolds*, Lect. Notes in Math., vol. 583, Springer Verlag, 1977.
- [44] E. Hopf, *Statistik der geodätischen Linien in Mannigfaltigkeiten negativer Krümmung*, Ber. Verh. Sächs. Akad. Wiss. Leipzig **91** (1939), 261–304.
- [45] V. Horita and A. Tahzibi, *Partial hyperbolicity for symplectic diffeomorphisms*, Ann. Inst. H. Poincaré Anal. Non Linéaire **23** (2006), 641–661.
- [46] A. Katok, *Lyapunov exponents, entropy and periodic points of diffeomorphisms*, Publ. Math. IHES **51** (1980), 137–173.
- [47] A. Katok and A. Kononenko, *Cocycle's stability for partially hyperbolic systems*, Math. Res. Lett. **3** (1996), 191–210.
- [48] F. Ledrappier, *Propriétés ergodiques des mesures de Sinai*, Publ. Math. I.H.E.S. **59** (1984), 163–188.
- [49] ———, *Positivity of the exponent for stationary sequences of matrices*, Lyapunov exponents (Bremen, 1984), Lect. Notes Math., vol. 1186, Springer, 1986, pp. 56–73.
- [50] F. Ledrappier and L.-S. Young, *The metric entropy of diffeomorphisms. I. Characterization of measures satisfying Pesin's entropy formula*, Ann. of Math. **122** (1985), 509–539.
- [51] ———, *The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension*, Ann. of Math. **122** (1985), 540–574.
- [52] C. Liverani, *On contact Anosov flows*, Ann. of Math. **159** (2004), 1275–1312.

-
- [53] J. Milnor, *Fubini foiled: Katok's paradoxical example in measure theory*, Math. Intelligencer **19** (1997), 30–32.
- [54] V. Nițică and A. Török, *An open dense set of stably ergodic diffeomorphisms in a neighborhood of a non-ergodic one*, Topology **40** (2001), 259–278.
- [55] Ya. Pesin, *Lectures on partial hyperbolicity and stable ergodicity*, European Mathematical Society (EMS), 2006, Zurich Lectures in Advanced Mathematics.
- [56] L. Polterovich, *Growth of maps, distortion in groups and symplectic geometry*, Invent. Math. **150** (2002), 655–686.
- [57] C. Pugh and M. Shub, *Ergodicity of Anosov actions*, Invent. Math. **15** (1972), 1–23.
- [58] ———, *Stable ergodicity and partial hyperbolicity*, International Conference on Dynamical Systems (Montevideo, 1995), Pitman Res. Notes Math. Ser., vol. 362, Longman, 1996, pp. 182–187.
- [59] ———, *Stably ergodic dynamical systems and partial hyperbolicity*, J. Complexity **13** (1997), 125–179.
- [60] ———, *Stable ergodicity and julienne quasi-conformality*, J. Europ. Math. Soc. **2** (2000), 1–52.
- [61] F. Rodríguez Hertz, *Stable ergodicity of certain linear automorphisms of the torus*, Ann. of Math. **162** (2005), 65–107.
- [62] F. Rodríguez Hertz, M. A. Rodríguez Hertz, and R. Ures, *A survey of partially hyperbolic dynamics*, Partially hyperbolic dynamics, laminations, and Teichmüller flow, Fields Inst. Commun., vol. 51, Amer. Math. Soc., 2007, pp. 35–87.
- [63] ———, *Accessibility and stable ergodicity for partially hyperbolic diffeomorphisms with 1D-center bundle*, Invent. Math. **172** (2008), 353–381.
- [64] ———, *Partial hyperbolicity and ergodicity in dimension three*, J. Mod. Dyn. **2** (2008), 187–208.
- [65] J. Rodríguez Hertz, M. A. Rodríguez Hertz, A. Tahzibi, and R. Ures, *A criterion for ergodicity of non-uniformly hyperbolic diffeomorphisms.*, www.arXiv.org arXiv:0710.2353.
- [66] D. Ruelle, *Perturbation theory for Lyapunov exponents of a toral map: extension of a result of Shub and Wilkinson*, Israel J. Math. **134** (2003), 345–361.
- [67] D. Ruelle and A. Wilkinson, *Absolutely singular dynamical foliations*, Comm. Math. Phys. **219** (2001), 481–487.
- [68] R. Saghin and Z. Xia, *Partial hyperbolicity or dense elliptic periodic points for c^1 -generic symplectic diffeomorphisms*, Trans. Amer. Math. Soc. **358** (2006), 5119–5138.
- [69] ———, *Geometric expansion, Lyapunov exponents and foliations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **26** (2009), 689–704.
- [70] M. Shub and A. Wilkinson, *Pathological foliations and removable zero exponents*, Invent. Math. **139** (2000), 495–508.
- [71] A. Tahzibi, *Stably ergodic diffeomorphisms which are not partially hyperbolic*, Israel J. Math. **142** (2004), 315–344.

-
- [72] M. Viana, *Almost all cocycles over any hyperbolic system have nonvanishing Lyapunov exponents*, Ann. of Math. **167** (2008), 643–680.
- [73] A. Wilkinson, *The cohomological equation for partially hyperbolic diffeomorphisms*, arXiv:0809.4862.
- [74] ———, *Stable ergodicity of the time-one map of a geodesic flow*, Ergod. Th. & Dynam. Sys. **18** (1998), 1545–1587.